



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Fetal surgery in zurich: key features of our first open in utero repair of myelomeningocele

Meuli, Martin ; Moehrlen, Ueli ; Flake, Alan ; Ochsenbein, Nicole ; Huesler, Margaret ; Biro, Peter ;
Scheer, Ianina ; Tharakan, Sasha ; Dürig, Peter ; Zimmermann, Roland

DOI: <https://doi.org/10.1055/s-0032-1329700>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-69574>

Journal Article

Originally published at:

Meuli, Martin; Moehrlen, Ueli; Flake, Alan; Ochsenbein, Nicole; Huesler, Margaret; Biro, Peter; Scheer, Ianina; Tharakan, Sasha; Dürig, Peter; Zimmermann, Roland (2013). Fetal surgery in zurich: key features of our first open in utero repair of myelomeningocele. *European Journal of Pediatric Surgery*, 23(06):494-498.

DOI: <https://doi.org/10.1055/s-0032-1329700>

Fetal Surgery in Zurich: Key Features of our First Open In Utero Repair of Myelomeningocele

Journal:	<i>European Journal of Pediatric Surgery</i>
Manuscript ID:	EJPS-2012-05-2152-CR
Manuscript Type:	Case Report
Date Submitted by the Author:	15-May-2012
Complete List of Authors:	<p>Meuli, Martin; Children's University Hospital Zurich, Department of Pediatric Surgery Moehrlen, Ueli; University Children's Hospital Zurich, Department of Surgery; Flake, Alan; The Children's Hospital of Philadelphia, and The University of Pennsylvania School of Medicine, The Center for Fetal Diagnosis and Treatment Ochsenbein, Nicole; University Hospital Zurich, Department of Obstetrics Huesler, Margaret; University Hospital Zurich, Department of Obstetrics Biro, Peter; University Hospital Zurich, Institute of Anesthesiology Scheer, Ianina; University Children's Hospital Zurich, Department of Radiology Tharakan, Sasha; Children's University Hospital Zurich, Department of Pediatric Surgery Dürig, Peter; Effinger Center for Obstetrics and Gynecology, - Zimmermann, Roland; University Hospital Zurich, Department of Obstetrics</p>
Keywords:	fetal surgery, myelomeningocele, MMC, open fetal surgery
Abstract:	<p>Background The recently published MOMS-Trial suggested that open in utero repair is a novel standard of care for select fetuses with MMC. At the Zurich Center for Fetal Diagnosis and Therapy we carried out one of the first fetal MMC repairs outside the three US-Trial Centers that was performed in strict accord with the recently published criteria, principles, and procedures set forth by the quoted trial.</p> <p>Case Report A 37 y-old 4G3P was diagnosed at GA 19+3 with a fetus demonstrating a lumbosacral MMC (upper level L3). Since all criteria for eventual fetal repair were met, the parents were non-directively counseled and then opted for fetal surgery. The operation was successfully performed at GA 24+5. The postoperative course of mother and fetus was uneventful except for leakage-induced oligohydramnios. The fetal middle cerebral artery peak velocity was moderately elevated for two weeks postoperatively suggesting fetal anemia. C-section was performed at GA 35 after premature rupture of membranes. At 8 months of life, a VP-shunt had to be placed. The peripheral neurology was determined as a mild to moderate paraparesis</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>(level L3/4). A bilateral hip dislocation due to prolonged breech presentation required surgery. There was no need for clean intermittent catheterization (CIC). Finally, the baby suffered from a kidney dysfunction of unclear origin.</p> <p>Conclusion</p> <p>We have adopted the novel and, more importantly, evidence-based treatment option resulting from the MOMS-trial and set the stage for a scientifically sound open fetal surgery program outside the US-Trial Centers. Maternal safety was perfectly preserved. Fetal problems possibly related to prenatal surgery comprised uneventful late preterm birth, oligohydramnios-related hip dislocations, and mild renal dysfunction. At one year of age, distinct benefits from fetal surgery include favorable leg motor function and avoidance of bladder catheterization.</p>

SCHOLARONE™
Manuscripts

Abstract

Background

The recently published MOMS-Trial suggested that open in utero repair is a novel standard of care for select fetuses with MMC. At the Zurich Center for Fetal Diagnosis and Therapy we carried out one of the first fetal MMC repairs outside the three US-Trial Centers that was performed in strict accord with the recently published criteria, principles, and procedures set forth by the quoted trial.

Case Report

A 37 y-old 4G3P was diagnosed at GA 19+3 with a fetus demonstrating a lumbosacral MMC (upper level L3). Since all criteria for eventual fetal repair were met, the parents were non-directively counseled and then opted for fetal surgery. The operation was successfully performed at GA 24+5. The postoperative course of mother and fetus was uneventful except for leakage-induced oligohydramnios. The fetal middle cerebral artery peak velocity was moderately elevated for two weeks postoperatively suggesting fetal anemia. C-section was performed at GA 35 after premature rupture of membranes. At 8 months of life, a VP-shunt had to be placed. The peripheral neurology was determined as a mild to moderate paraparesis (level L3/4). A bilateral hip dislocation due to prolonged breech presentation required surgery. There was no need for clean intermittent catheterization (CIC). Finally, the baby suffered from a kidney dysfunction of unclear origin.

Conclusion

We have adopted the novel and, more importantly, evidence-based treatment option resulting from the MOMS-trial and set the stage for a scientifically sound open fetal surgery program outside the US-Trial Centers. Maternal safety was perfectly preserved. Fetal problems possibly related to prenatal surgery comprised uneventful late preterm birth, oligohydramnios-related hip dislocations, and mild renal dysfunction. At one year of age, distinct benefits from fetal surgery include favorable leg motor function and avoidance of bladder catheterization.

Introduction

The first description of apparently secondary, i.e. in utero acquired, damage to the pathologically exposed spinal cord within the myelomeningocele (MMC) lesion dates back to 1956 (1). For obvious reasons, the significance of this observation regarding therapeutic consequences was not recognized until fetal surgery became a reality in the 1980ies (2). Only then was the hypothesis born that early in utero intervention might stop the ongoing neural tissue destruction and so reduce the neurologic deficit otherwise seen at birth (3).

Compelling experimental (4; 5) as well as preliminary clinical (6-10) evidence collected over the past two decades have led to the conclusion that fetal surgery for MMC may be an effective novel therapy to ameliorate the neurologic outcome. Lastly, a recently published prospective, controlled, and randomized study (MOMS-Trial) (11) has, for the very first time, generated sound data showing that in utero repair can be recommended as a novel standard of care for selected fetuses with MMC. This is a report on the first fetus with MMC operated under the published criteria, principles, and procedures set forth by the MOMS-Trial at the Zurich Center for Fetal Diagnosis and Therapy.

Case report

Diagnostic workup

A healthy 37 year-old 4G3P was diagnosed with a female fetus suffering from a spina bifida in the 19th week of gestation. On ultrasound, the fetus had a classical cystic lumbosacral myelomeningocle (MMC) (cranial level = lumbar 3), the typical Chiari II malformation, and a moderate ventriculomegaly (13 mm). The fetus demonstrated bilateral lower extremity movements at hip, knee, ankle, and toe levels. The bladder showed normal cycling and the stomach was visible. No other malformations were identified. The placenta completely covered the anterior uterine wall. Fetal MRI at 19+4 weeks of gestational age confirmed these findings, but additionally revealed an aqueductal stenosis (Fig. 1a).

In summary, both mother and fetus met all inclusion criteria for fetal surgical repair according to the MOMS-Trial while they did not fulfill any of the exclusion criteria (11).

Prenatal counseling

Parents were counseled non-directively in several sessions by pediatric surgeons, obstetricians, and anesthesiologists with particular focus on all known risks and potential benefits associated with maternal-fetal surgery and also with all problems and handicaps presumably present if standard postnatal care was chosen. They were given ample time for decision making and were also informed that whatever decision they made they could change at any time.

The parents decided to pursue fetal surgery. A written informed consent was obtained.

Maternal-fetal surgery

Both maternal and fetal surgery were performed according to the current techniques used within the MOMS-Trial framework (11). Three days before the intervention lung maturation was induced with 24mg betamethasone. To prevent postoperative contractions, the mother received 50mg indomethacin preoperatively. The mother underwent deep inhalation anesthesia with desflurane, remifentanyl and neuromuscular relaxation. A transverse supraumbilical laparotomy was performed, the uterus was exteriorized, and opened on the posterior wall at a safe distance from the placenta. The fetus was positioned in such a way that the cystic MMC-lesion came to lie within the hysterotomy.

The cystic sac was then resected taking care not to touch the exposed, non-neurulated, but macroscopically intact, spinal cord tissue (Fig. 2a). The easily identifiable filum terminale was clearly under tension that was completely released after untethering. Since the open dura mater was difficult to identify, a bilateral paraspinal myofascial flap, medially comprising the open dura, was raised and sewn over the cord (Fig 2b). After extensive mobilization, the skin could be closed primarily over the defect (Fig 2c). Total fetal blood loss was an estimated 8 ml (~ 8-10% loss of total blood volume).

Ultrasound monitoring showed normal heart rate and myocardial contractility of the fetus throughout the intervention.

The uterus was closed using an inner running suture and outer single stitches. Amniotic fluid was substituted with 500 ml Ringer's lactate containing 2 g of ceftriaxon as antibiotic prophylaxis. An omental flap was sewn over the closure site for additional sealing. The maternal laparotomy was closed in layers.

In summary, the operation was performed without significant problems and the principle goals of the intervention including preserved maternal-fetal safety as well as solid multilayer defect closure over the non-touched intact spinal cord were reached.

Postoperative maternal-fetal management and evolution

To prevent contractions, an infusion with high dose magnesium was started (6 g loading dose, followed by 4 g/h for the first 3 h, then 3 g/h for the next 33 h) and the mother also received 3 doses of 50 mg indomethacin. After 3 days, a continuous long-term tocolysis was started using nifedipine 120 mg/d. For thromboprophylaxis we used low molecular heparin and pneumatic stockings. Epidural catheter analgesia allowed optimal pain control for the first 2 postoperative days. The mother was mobilized on postoperative day 2 and discharged home 4 weeks after the intervention.

Continuous monitoring of fetal heart rate and uterine activity was normal throughout the postoperative course.

A scan 6 h after the intervention demonstrated anhydramnios suggesting amniotic fluid leakage into the maternal peritoneal cavity although no free fluid was seen in the maternal abdomen. For the remainder of gestation, the fetus remained in the breech presentation with extended legs, already present preoperatively.

Postoperatively, peak velocity in the middle cerebral artery (MCA-PV) was 2.7 SD above mean values, indicating mild fetal anemia. It increased to 3.7 SD within one week and normalized by two weeks after the operation. Heart rate and myocardial contractility were always normal and there were no signs of fetal hydrops. The fetal bladder was almost empty for one week. Thereafter, normal cycling was again present and small pockets of amniotic fluid were detected. The amniotic fluid index remained below 5. The ventricle size was 11 mm during the first 2 postoperative weeks but then gradually increased to 20mm prior to delivery. Hindbrain herniation reversed completely within 8 weeks (Fig 1b). Active feet and toe movements were present throughout.

Cesarean section

C-section was performed on a semi-elective basis at gestational age of 35+0 weeks, about 8 hours after rupture of membranes and onset of mild labour. There was no uterine dehiscence detectable.

The baby girl (head circumference 32,5 cm [P:50-75], weight 2030 g [P:10-25], length 43 cm [P:3-10]) adapted normally (Apgar 8/8/9). The repair site was solidly healed (Fig. 3). Due to prolonged breech presentation, legs and trunk were in a fixed jack-knife position. The mother exhibited an uneventful postoperative course and was discharged home on postoperative day 6.

Postnatal clinical evolution during the first year of life

The key features of the first 12 months post partum can be summarized as follows: Vital parameters were always normal. Regarding growth, the baby developed along the 3rd percentile for length and body weight. Peripheral neurology showed a mild to moderate paraparesis below L4 representing a more favourable motor capacity than usually expected when compared to similar postnatal repair.

With regard to hydrocephalus, head circumference increased gradually from the percentile P3 (birth) to P50 (6 months) without clinically significant signs of increased cerebral pressure. Over the ensuing 2 months, the baby demonstrated an overproportional growth of the neurocranium, developed mild signs for increased cerebral pressure, and head circumference crossed P90. Repeat MRI demonstrated significant ventriculomegaly (Fig 4), in all likelihood due to the still present aqueductal stenosis (present already before fetal

surgery!), while hindbrain herniation was still reversed. At 8 months of life, a ventriculo-peritoneal shunt was therefore inserted with an uneventful postoperative course up to date.

Urologically, the baby is voiding spontaneously under a regimen of oxybutinin installation twice a day.

The baby demonstrated a renal insufficiency with the highest creatinin level of 273 $\mu\text{mol/l}$ at day 5 of life that gradually decreased to 48 $\mu\text{mol/l}$ (normal range $<35 \mu\text{mol/l}$) at the age of 11 months. Despite an extensive diagnostic workup, the reason for this kidney dysfunction could not be conclusively determined.

Defecation occurs spontaneously and on a regular basis without soiling.

Finally, a bilateral hip dislocation, consequent to the fetal breech position (present before fetal surgery) was reduced surgically at 4 months of life after casting failed.

Discussion

This is one of the first open fetal spina bifida repairs conducted outside the USA. More importantly, it is, to the best of our knowledge, the first intervention performed outside the three US-trial centers *after* the ground breaking evidence resulting from the “Management of Myelomeningocele Study” (MOMS) clinical trial was published (11). Of note, this “premiere” surgery was carefully orchestrated with our partner institution, the Center for Fetal Diagnosis and Treatment of the Children’s Hospital of Philadelphia. In particular, the current concepts, guidelines, and standard operating procedures of the MOMS-Trial regarding prenatal diagnostic workup, maternal-fetal surgery, and postoperative management were adopted (11). Importantly, one of the world’s most experienced fetal surgeons (co-author AWF, Philadelphia) assisted the operation and acted as advisor during the critical perioperative phase. It is our firm conviction that the latter is an indispensable strategic element to guarantee the best possible setting for this most sophisticated and extremely demanding enterprise. Clearly, a failure to organize the best expertise is hardly justifiable medically and ethically in front of patient and mother, and, moreover, it jeopardizes the successful spawn of this youngest surgical discipline to a few more select centers.

Definitely, the aforementioned MOMS-Trial is by far the largest prospective, randomized clinical trial ever carried out to compare prenatal versus postnatal surgery. And it is the first to convincingly demonstrate a clear cut benefit for the fetal surgery cohort in that almost all crucial outcome parameters were clearly in favor of in utero intervention as opposed to conventional postnatal care (11). Of note, the unequivocal results even led to a premature ending of the trial after 183 out of the planned 200 patients were enrolled.

The key findings of the MOMS-Trial were that fetal surgery markedly reduced the need for hydrocephalus shunting (fetal surgery 40% vs. postnatal care 82%) and significantly improved motor as well as developmental outcomes at 30 months. On the other hand, fetal surgery was associated with a higher risk of preterm delivery and uterine dehiscence at delivery (11).

We found similar advantageous effects in our case, in particular complete reversal of hindbrain herniation, spontaneous micturition with no need for clean intermittent catheterization, and a better than expected peripheral neurology. Although these findings are substantial benefits attributable in all likelihood to prenatal intervention, they are somewhat preliminary since the follow-up period is only one year.

The need for late ventriculo-peritoneal shunting at 8 months of life calls for a judicious consideration. The hindbrain herniation, present preoperatively, resolved almost completely before birth as typically seen after fetal surgery (11-15). During the first 6 months there was a slightly overproportional head growth and there were no signs for increased intracranial pressure (ICP). Therefore, we expected a high likelihood that shunting would not be mandatory. Somewhat surprisingly, we then witnessed an unfavorable evolution with increasing signs for elevated ICP and a head circumference that rapidly crossed the 97th percentile mandating shunt placement. Here, the question arises whether shunting was required because of aqueductal stenosis, identified before fetal surgery. We did not find any current literature describing an association between MMC and aqueductal stenosis. Therefore, it remains unclear whether this patient is a true non-responder regarding shunt avoidance after fetal surgery (like the 40% of the MOMS Trial) or whether it was a rather unfortunate and uncommon coincidence of spina bifida and aqueductal stenosis leading to aqueductal stenosis-induced/co-induced hydrocephalus (16).

Another issue is the leakage-induced oligohydramnios (also seen in 21% of the MOMS-Trial fetal surgery cases) and its possible negative impact. Apparently, there was no clinically significant effect on lung development and function in that the baby demonstrated normal respiration at all times. Likewise, oligohydramnios has probably not played a major role in the pathogenesis of hip dislocation. Rather, this pathology is a consequence of the breech presentation the fetus demonstrated already *before* fetal intervention.

A completely unexpected finding was a marked transient renal insufficiency of unclear origin. We hypothesize that a combined effect of indomethacin (17; 18) (via renal artery constriction) and anemia, another drug-related side effect, or a non procedure-related cause (e.g. hypodysplastic kidneys) might have played a role. Despite extensive investigation we could not find enough evidence to corroborate either explication, although a pharmacological problem is most likely. Moreover, there is no report on procedure-related kidney problems in the context of fetal surgery.

Of utmost importance, maternal safety was perfectly preserved as there were no significant problems or complications throughout the entire course.

In summary, we report on our first case successfully managed in accord with the recently published MOMS-Trial framework and provide preliminary outcome data. Although in

1
2
3 utero surgery can now be recommended as new standard of care for select fetuses with
4 MMC, the paucity of cases as well as both delicacy and intricacy of fetal surgery mandate
5 these patients be pooled in a few centers worldwide and the interventions be exclusively
6 performed by appropriately trained and highly qualified fetal surgeons.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Reference List

- 1 Cameron AH. The spinal cord lesion in spina bifida cystica. *Lancet* 1956;271:171-174.
- 2 Harrison MR, Filly RA, Golbus MS et al. Fetal treatment 1982. *N Engl J Med* 1982;307:1651-1652.
- 3 Hutchins GM, Meuli M, Meuli-Simmen C et al. Acquired spinal cord injury in human fetuses with myelomeningocele. *Pediatr Pathol Lab Med* 1996;16:701-712.
- 4 Meuli M, Meuli-Simmen C, Hutchins GM et al. In utero surgery rescues neurological function at birth in sheep with spina bifida. *Nat Med* 1995;1:342-347.
- 5 Stiefel D, Copp AJ, Meuli M. Fetal spina bifida in a mouse model: loss of neural function in utero. *J Neurosurg* 2007;106:213-221.
- 6 Adzick NS, Sutton LN, Crombleholme TM et al. Successful fetal surgery for spina bifida. *Lancet* 1998;352:1675-1676.
- 7 Meuli M, Meuli-Simmen C, Yingling CD et al. Creation of myelomeningocele in utero: a model of functional damage from spinal cord exposure in fetal sheep. *J Pediatr Surg* 1995;30:1028-1032.
- 8 Meuli M, Meuli-Simmen C, Yingling CD et al. In utero repair of experimental myelomeningocele saves neurological function at birth. *J Pediatr Surg* 1996;31:397-402.
- 9 Meuli M, Meuli-Simmen C, Hutchins GM et al. The spinal cord lesion in human fetuses with myelomeningocele: implications for fetal surgery. *J Pediatr Surg* 1997;32:448-452.
- 10 Bruner JP, Tulipan N, Paschall RL et al. Fetal surgery for myelomeningocele and the incidence of shunt-dependent hydrocephalus. *JAMA* 1999;282:1819-1825.
- 11 Adzick NS, Thom EA, Spong CY et al. A randomized trial of prenatal versus postnatal repair of myelomeningocele. *N Engl J Med* 2011;364:993-1004.
- 12 Bouchard S, Davey MG, Rintoul NE et al. Correction of hindbrain herniation and anatomy of the vermis after in utero repair of myelomeningocele in sheep. *J Pediatr Surg* 2003;38:451-458.
- 13 Sutton LN, Adzick NS, Bilaniuk LT et al. Improvement in hindbrain herniation demonstrated by serial fetal magnetic resonance imaging following fetal surgery for myelomeningocele. *JAMA* 1999;282:1826-1831.
- 14 Tulipan N, Hernanz-Schulman M, Bruner JP. Reduced hindbrain herniation after intrauterine myelomeningocele repair: A report of four cases. *Pediatr Neurosurg* 1998;29:274-278.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 15 Tulipan N, Hernanz-Schulman M, Lowe LH et al. Intrauterine myelomeningocele repair reverses preexisting hindbrain herniation. *Pediatr Neurosurg* 1999;31:137-142.
- 16 Jellinger G. Anatomopathology of non-tumoral aqueductal stenosis. *J Neurosurg Sci* 1986;30:1-16.
- 17 Gloor JM, Muchant DG, Norling LL. Prenatal maternal indomethacin use resulting in prolonged neonatal renal insufficiency. *J Perinatol* 1993;13:425-427.
- 18 van der Heijden BJ, Carlus C, Narcy F et al. Persistent anuria, neonatal death, and renal microcystic lesions after prenatal exposure to indomethacin. *Am J Obstet Gynecol* 1994;171:617-623.




Figure 1

Figure 1




Figure 2

Figure 2



Figure 3

Figure 3

Peer Review




Figure 4

Figure 4

Legend for figures:

Figure 1

a) preoperative fetal MRI at 20 weeks of gestation. Sagittal single shot fast spin echo sequence showing the MMC lesion (asterisk) with associated hindbrain herniation (white arrow) and ventriculomegaly.
b) postoperative fetal MRI at 34 weeks of gestation. Sagittal single shot fast spin echo sequence showing an intact MMC closure side. The hindbrain herniation is reversed. Dysplastic mesencephalic tectum.

Figure 2

a) MMC lesion
b) Closure of the dura mater
c) fetal skin closure

Figure 3

Picture taken immediately after birth showing a healed wound and covered spina bifida.

Figure 4

MRI at the 8 month of age; Sagittal T2-weighted fast spin echo sequence: Small posterior fossa, no cerebellar herniation. Dysplastic mesencephalic tectum and missing flow void at the level of the cerebral aqueduct (arrow).